

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 108 364 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
20.06.2001 Bulletin 2001/25

(51) Int Cl.7: **A23D 9/007**, **A23L 1/24**,
A23L 1/30

(21) Application number: **00311172.1**

(22) Date of filing: **14.12.2000**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• **Dartey, Clemence K.**
North Wales, PA 19545 (US)
• **Sox, Thomas E.**
Ambler, PA 19002 (US)

(30) Priority: **15.12.1999 US 461607**

(74) Representative: **Mercer, Christopher Paul**
Carpmaels & Ransford
43, Bloomsbury Square
London WC1A 2RA (GB)

(71) Applicant: **McNEIL-PPC, INC.**
Skillman, NJ 08858 (US)

(54) Long chain alcohols admixed in sterol compounds

(57) The present invention provides a method of providing a desirable admixture of long chain alcohols and sterols in a food or beverage product. More specif-

ically, this process comprises dissolving or suspending the long chain alcohol in the sterol, and then incorporating the admixture into food or beverage product.

EP 1 108 364 A2

Description**Related Applications**

- 5 [0001] This application is being concurrently filed with two additional applications. These related applications are entitled Long Chain Alcohols Provided in Edible Oils, European Patent Application No. , claiming priority from USSN 09/461 887 (Attorney's ref: P026182EP) European Patent Application No. , claiming priority from USSN 09/461 592 (Attorney's ref: P026184EP).

Field of the Invention

- 10 [0002] The present invention relates to the incorporation of long chain alcohol substances, more specifically the present invention relates to the encapsulation of long chain alcohols so as to more readily incorporate them into foods.

Background of the Invention

- 15 [0003] Long chain alcohols, including policosanols, are therapeutically useful materials with efficacy in lowering cholesterol or blood lipids, inhibiting platelet aggregation and improving stamina. Therefore, the incorporation of long chain alcohols such as policosanols in various foods is desirable.
- 20 [0004] Incorporation of policosanols into high fat or fat-continuous emulsion systems such as margarine and margarine spreads is complicated by the functional properties of policosanols. In particular, incorporation of policosanols into a margarine oil system containing diglycerides and a phospholipid causes an increase in the hardness of a margarine type product as disclosed in WO 98/47385. In addition, EP 0991804 discloses that the incorporation of a natural long chain alcohol in a fat continuous system reduces the viscosity and yield values of confectionery products.
- 25 [0005] Despite these disclosures there is still an ongoing need to be able to add long chain alcohols to foods in a manner that the long chain alcohol will be easily formulated, will remain stable during storage, and will not adversely affect the properties of the food.

Summary of the Invention

- 30 [0006] The present invention provides comestibles that contain an admixture of long chain alcohols and sterol compounds. Both the long chain alcohols and sterol compounds are provided in the amounts effective to lower cholesterol levels in a human being when consumed on a regular basis.

Detailed Description of the Invention

- 35 [0007] The present invention is directed to the admixture of long chain alcohols into sterol compounds and the incorporation of the alcohol/sterol mixtures into food or pharmaceutical products. By long chain alcohols, it is understood to mean both saturated and unsaturated alcohols of C₂₀ and above, primarily from C₂₀ to C₃₆. As used herein, long chain alcohols are understood to be those materials that contain more than about 90 weight percent C₂₀ or longer, primarily aliphatic alcohol materials. For the greatest health benefit it is preferred that the long chain alcohols be more than 50 weight percent octacosanol (C₂₈), preferably more than 65 percent, more preferably greater than about 70 weight percent. As used herein policosanols are understood to be a mixture of long chain alcohols ranging from C₂₀ to C₃₆ with greater than 50 weight percent C₂₈, preferably greater than 65 weight percent octacosanol. Common distribution and concentration ranges of the various components of policosanols are disclosed in US Patent 5,856,316, hereby incorporated by reference. These long chain alcohols are available from various natural sources, most preferably from sugar cane wax. The long chain alcohols can also be synthesized using techniques well known in the art.
- 40 [0008] The level of long chain alcohols incorporated in a food is determined by serving size of the food product. It is desirable to provide in a single serving an effective amount of the long chain alcohols to derive the desired physiological benefits, such as reduced platelet aggregation, reduced lipid and cholesterol levels and the like. Typically the level of the long chain alcohols is from about 0.5 to about 400 mg/serving size; preferably from about 1.0 to about 100 mg/serving and most preferably from about 2.0 to about 20 mg/serving size of a food product. When used in pharmaceutical dosage forms, it is preferred that the level of the long chain alcohol is less than about 40 milligrams, preferably from about 2 to about 20 milligrams so as to maintain a stable dosage form.
- 45 [0009] There are many foods into which the encapsulated long chain alcohols are incorporated, including but not limited to margarine, spreads, salad dressings, beverages and juices, cookies, confectionery products, creams, cheeses, oils, candy and the like.
- 50 [0010] The sterol materials in which the long chain alcohols are dissolved are well known in the art. As used herein

the term sterol is understood to include cholesterol-lowering materials including sterols, stanols, stanol esters, sterol esters. Among the preferred materials are β -sitosterol and β -sitostanol, as well as oryzanol and mixtures of various sterols and stanols. These materials are well known in the art and are disclosed in U.S. Patent Nos. 3,881,005; 5,502,045; 5,869,708, hereby incorporated by reference and WO 98/19556. The level of sterols used in the present invention should be sufficient to provide an amount effective in lowering cholesterol in a vertebrate, such as mammals, reptiles and amphibians, preferably a human when consumed on a routine basis. Typically the sterol level is from about 0.4 to about 4.0 g, preferably from about 0.8 to about 3.0 g, and most preferably from about 1.0 to about 2.0 g per serving.

[0011] The relative weight ratios of the long chain alcohols and the sterol compounds are such that the long chain alcohols will be admixed into the sterol compounds. In a preferred embodiment, the admixture is then heated to melt completely before adding the liquid blend to the oil phase or the aqueous phase compositions. In preparing a cholesterol-lowering margarine or a margarine spread, the long chain alcohols are mixed with melted stanol esters at about 140 °F. The mixture is heated to 180 °F to dissolve completely the long chain alcohols. The hot alcohol/stanol ester preparation is added to the margarine oil stabilized at 160 °F before the aqueous phase is mixed with the oil phase to produce the margarine emulsion. In a preferred embodiment of this invention, the mixture of sterol compounds and long chain alcohols are heated and dissolved in a portion of the margarine oil at about 180 °F. This oil blend is then added to remaining oil phase maintained at 160 °F prior to the addition of the aqueous phase to form the margarine emulsion. At temperatures above 140 °F, policosanols preparations, as high as 1.0 %, remain dissolved in stanol esters, sterol esters, sterols, stanols and the like.

[0012] Emulsifying agents and surfactants can also be employed to aid in admixing the long chain alcohols in the sterol compounds. Suitable emulsifying agents and suspending agents include any food grade materials, such as mono and diglycerides, TWEENS and SPANS, lecithin, polyglycerol esters, propylene glycol esters and the like, polysorbates, mono and diglycerides of fatty acids, sucrose fatty acid esters and polyoxyethylene derivatives of sorbitan fatty acid esters. These surfactants are well known in the art and are commercially available.

[0013] Suitable polyglycerol esters include triglyceryl monostearate, hexaglyceryl distearate, hexaglyceryl monopalmitate, hexaglyceryl dipalmitate, decaglyceryl distearate, decaglyceryl monooleate, decaglyceryl dioleate, decaglycerol monopalmitate, decaglycerol dipalmitate, decaglyceryl monostearate, octaglycerol monooleate, octaglycerol monostearate and decaglycerol monocaprylate.

[0014] Other useful surfactants include polysorbates made from the reaction product of monoglycerides or sorbitan esters with ethylene oxides. Examples of useful polysorbates include polyoxyethylene 20 mono- and diglycerides of saturated fatty acids, polyoxyethylene 4 sorbitan monostearate, polyoxyethylene 20 sorbitan tristearate, polyoxyethylene 20 sorbitan monooleate, polyoxyethylene 5 sorbitan monooleate, polyoxyethylene 20, sorbitan trioleate, sorbitan monopalmitate, sorbitan monolaurate, propylene glycol monolaurate, glycerol monostearate, diglycerol monostearate, glycerol lactyl-palmitate.

[0015] Other suitable surfactants with HLB values provided in brackets, [], include decaglycerol monolaurate [15.5]; decaglycerol distearate [10.5]; decaglycerol dioleate [10.5]; decaglycerol dipalmitate [11.0]; decaglycerol monostearate [13.0]; decaglycerol monooleate [13.5]; hexaglycerol monostearate [12.0]; hexaglycerol monooleate [10.5]; hexaglycerol monoshortening [12.0]; polyoxyethylene (20) sorbitan monolaurate [16.7]; polyoxyethylene (4) sorbitan monolaurate [13.3]; polyoxyethylene (20) sorbitan monopalmitate [15.6]; polyoxyethylene (20) sorbitan monostearate [14.9]; polyoxyethylene (20) sorbitan tristearate [10.5]; polyoxyethylene (20) sorbitan monooleate [15.0]; polyoxyethylene (5) sorbitan monooleate [10.0]; polyoxyethylene (20) sorbitan trioleate [11.0]. As is appreciated by those with skill in the art, the HLB value for a surfactant is an expression of its Hydrophile-Lipophile Balance, i.e., the balance of the size and strength of the hydrophilic (polar) and lipophilic (non-polar) groups of the surfactant.

[0016] Lactic acid derivatives include sodium stearyl lactylate and calcium stearyl lactylate. Most preferably the emulsifying materials are polysorbates and the partial glycerides, including mono and diglycerides.

[0017] Another technique which can be employed is to melt and dissolve both the long chain alcohol and sterol compound in a suitable solvent. Any suitable food-grade vegetable fats and oils can be employed as solvents for the long chain alcohols and the sterol compounds. Aqueous based systems can also be employed, but this will require the use of various emulsifying agents approved for food applications as set forth above.

[0018] In a highly preferred embodiment, other ingredients which have desirable physiological actions can be added to the food products. Such materials include soy, vitamins, minerals and the like. Also included in the comestible are other ingredients typically used in the preparation of foods which include without limitation, flavors, colors, preservatives, sweeteners and the like.

[0019] US Patent No. 5,952,393 (Sorkin) discloses tablet and soft gelatin compositions containing an impure policosanols obtained from rice bran and sterol blend. In contrast, the present invention is directed to the incorporation of an intimate admixture of long chain alcohol and phytosterols into a food or beverage product. Sterols are believed to lower serum cholesterol levels by blocking the absorption of dietary cholesterol and preventing the resorption of bile acids secreted during the digestive process. Therefore sterols are most efficacious when incorporated in the form of a food or beverage product. When incorporated into a food or beverage, the sterols are intimately mixed with the other

ingredients of the food into which they are formulated. Additionally the process of mastication and gastric mixing provides mixing of the sterols with cholesterol containing food that may be ingested in the same meal.

[0020] Sorkin also discloses the use of from about 58 to 83 milligrams of rice bran wax derived policosanols per tablet or soft gelatin capsule. In contrast, the present invention is directed to the administration of about 2 to about 20 milligrams of long chain alcohol, preferably policosanols per serving. The lower levels of the long chain alcohols are still therapeutically effective, particularly when consumed in combination with sterol compounds. Another advantage of the present invention is that the long chain alcohol/sterol combination is stable. Stable, as used herein means that the long chain alcohol does not precipitate or separate from the sterol compound.

[0021] The following examples are provided as specific embodiments of the present invention. Other modifications of this invention will be readily apparent to those skilled in the arts without departing from the scope of the present invention. The source of the material used in an example is provided in parenthesis.

Example 1

[0022] This example discloses the application of policosanols with stanol esters in a margarine spread formulation. The mixture of policosanols and the sterol derivative is stirred and heated in a hot water bath to 180 °F. The clear policosanols preparation is added to the oil phase components of the cholesterol-lowering margarine spread.

Table 1:

Cholesterol Lowering Margarine Spreads Containing Policosanols Dissolved in Sterol Derivatives				
Ingredients	Stanol Esters Weight (lbs.)	sterol Esters weight (lbs.)	tanols weight (lbs.)	terols weight (lbs.)
Liquid Soybean Oil	11.7923	1.7923	1.7923	1.7923
Partially Hydrogenated Soybean Oil	11.0000	1.0000	5.0000	5.0000
Canola Oil	30.0000	0.0000	5.0510	5.0510
Plant Stanol Esters	21.5500	1.5500	2.4990	2.4990
Plant Sterol Esters	-			
Plant Stanols	-			
Plant Sterols	-			
Monoglycerides	0.3000	.3000	.3000	.3000
Lecithin	0.2000	.2000	.2000	.2000
Hexaglycerol Distearate, Polyaldol 6-2-S	0.1000	.1000	.1000	.1000
Artificial Butter Flavor	0.0500	.0500	.0500	.0500
Vitamins A and D	0.00625	.00625	.00625	.00625
Beta Carotene 30 % Solution	0.00150	.00150	.00150	.00150
Policosanols, 93.16 % Active, Garuda International, CHOLESTANOL	0.0679	.0679	.0679	.0679
Aqueous Phase				
Water	22.8175	2.8175	2.8175	2.8175
Salt	2.0000	.0000	.0000	.0000
Citric Acid	0.0075	.0075	.0075	.0075

Table 1: (continued)

Cholesterol Lowering Margarine Spreads Containing Policosanol Dissolved in Sterol Derivatives				
Ingredients	Stanol Esters Weight (lbs.)	terol Esters eight (lbs.)	tanols eight lbs.)	terols eight lbs.)
Calcium Disodium EDTA	0.0070	.0070	.0070	.0070
Potassium Sorbate	0.1000	.1000	.1000	.1000
Total	100.000	00.000	00.000	00.000

[0023] The margarine emulsion was prepared in a pilot plant by blending the aqueous phase with the oil phase. The oil phase was prepared by blending together the liquid soybean oil, partially hydrogenated soybean oil, canola oil, monoglycerides, lecithin and hexaglycerol distearate in a jacketed margarine emulsion tank. The tank was heated with hot water to 160 °F to melt and produce a clear liquid oil blend.

[0024] Policosanol was mixed with stanol esters in a stainless steel container. The container was placed in a water-bath on a stove and heated to approximately 180 °F to melt. The clear liquid policosanol preparation was transferred and mixed with the oil blend in the emulsion tank. The butter flavor, vitamins A and D blend and beta-carotene color were blended with the oil to complete the oil phase component of the margarine spread. The aqueous phase was prepared by dissolving salt, citric acid, ethylenediaminetetra acetic acid (EDTA) and potassium sorbate in the water. To prepare the margarine emulsion, the emulsion tank agitator was turned on at high speed and the aqueous phase was blended slowly into the oil phase to form the margarine emulsion. The temperature of the margarine emulsion was adjusted to 130 ± 5 °F. The margarine spread emulsion was pumped and chilled in a scrape surface heat exchanger, through to a pin worker to a filling unit. The temperature of the margarine spread at the filling unit was maintained at 45 - 65 °F by adjusting the temperature of the emulsion in the scrape surface heat exchanger. The finished margarine was packed in suitable containers and stored refrigerated. Each margarine formulation contained approximately 5 mg policosanol per 8 gram serving size.

Example 2

[0025] This example discloses cholesterol-lowering light margarine spreads containing policosanol dissolved in sterol derivatives.

Table 2:

Cholesterol-Lowering Light Margarine Spreads Containing Policosanol Dissolved in Sterol Derivatives				
Ingredients	tanol Esters	Sterol Esters %	Stanols %	Sterols %
Liquid Soybean Oil	7.9913	7.9913	37.0423	37.0423
Plant Stanol Esters	1.5500	1.5500	-	-
Plant Sterol Esters			-	-
Plant Stanols			12.4990	-
Plant Sterols			-	12.4990
Mono-glycerides	.4000	.4000	.4000	.4000
Lecithin	.3000	.3000	.3000	.3000
Hexaglycerol Distearate, Polyaldo 6-2-S	.1000	.1000	.1000	.1000
Hexaglycerol Mixed Esters, CAPROL ET	.1000	.1000	.1000	.1000
Artificial Butter Flavor	.0500	.0500	.0500	.0500
Vitamins A and D	.00625	.00625	.00625	.00625
Beta Carotene 30 % Solution	.00250	.00250	.00250	.00250
Policosanol, 93.16 % Active, Garuda CHOLESTANOL	.0679	.0679	.0679	.0679
Aqueous Phase				

Table 2: (continued)

Cholesterol-Lowering Light Margarine Spreads Containing Policosanol Dissolved in Sterol Derivatives				
Ingredients	stanol Esters	Sterol Esters %	Stanols %	Sterols %
Water	7.3150	7.3150	7.3150	7.3150
Salt	.0000	.0000	.0000	.0000
Potassium Sorbate	.1000	.1000	.1000	.1000
Citric Acid	.0100	.0100	.0100	.0100
Calcium Disodium EDTA	.0070	.0070	.0070	.0070
Total	00.000	00.000	00.000	00.000

[0026] Liquid soybean oil, partially hydrogenated vegetable oil, monoglycerides, lecithin, hexaglycerol distearate and hexaglycerol mixed esters were mixed together in a margarine emulsion tank and heated to 160 °F. Policosanol and stanol esters were mixed together in a stainless steel container and heated in a hot water bath to 180 °F. The melted policosanol preparation was blended into the oil in the emulsion tank. The butter flavor, Vitamins A & D blend and beta-carotene color were mixed into the oil to make up the oil phase. The water phase was prepared by dissolving the salt, potassium sorbate, citric acid and EDTA in the water. While mixing the oil phase at high speed, the aqueous phase was added slowly to form the margarine spread emulsion. The margarine emulsion was processed into a finished margarine spread by processing the emulsion through a scrape surface heat exchanger, through a pin worker or a B-unit through to a filler unit. The temperature of the chilling unit (the scrape surface heat exchanger) was regulated to produce a finished light margarine spread with a temperature of approximately 45 - 65 °F at the filler unit. The finished light margarine spreads were packed in suitable containers and stored refrigerated. Each light margarine spread formulation shown in Table 2 contained approximately 5 mg policosanol per 8 gram serving size.

Example 3

[0027] This example concerns the use of stanol esters (and or sterol esters) as a carrier system for policosanol application in dressings. Policosanol was heated to dissolve in liquid stanol esters; or policosanol was mixed with solid stanol esters and heated to melt and form a clear liquid. The liquid policosanol preparation was then used to produce cholesterol lowering dressings e.g., ranch, French, creamy Italian, thousand island and mayonnaise dressings.

Table 3:

Cholesterol-Lowering Ranch Dressings Containing (a) Policosanol and Stanol Esters and (b) Policosanol and Sterol Esters		
	(a)	(b)
	Stanol Esters	Sterol Esters
Ingredients	Weight (g)	Weight (g)
Soybean Oil, Salad Oil	480.00	480.00
Vinegar, 120 Grain, White distilled	84.00	84.00
Stanol Esters	71.34	-
Sterol Esters	-	71.34
Sugar, Fine Granulated	58.80	58.80
Ranch Seasoning	50.40	50.40
Cultured Buttermilk Solids, Armour	19.20	19.20
Salt	8.40	8.40
CAPROL ET, A. C. Humko	7.44	7.44
Egg Yolk, Liquid, NEPA #2, 10 % Salt	4.32	4.32
KELTROL T Xanthan Gum, Kelco	4.20	4.20

Table 3: (continued)

Cholesterol-Lowering Ranch Dressings Containing (a) Policosanol and Stanol Esters and (b) Policosanol and Sterol Esters		
	(a)	(b)
	Stanol Esters	Sterol Esters
Ingredients	Weight (g)	Weight (g)
Polysorbate 60	3.60	3.60
Lemon Juice Concentrate, 400 GPL	3.00	3.00
Propylene Glycol Alginate, KELCOLOID LVF, Kelco	2.10	2.10
Polysorbate 80, TWEEN 80	1.92	1.92
Titanium Dioxide	1.68	1.68
Potassium Sorbate	0.96	0.96
Sodium Benzoate	0.96	0.96
Vitamin E Acetate Roche	0.22	0.22
Policosanol, 95 % Active, Garuda Cholesstanol	0.42	0.42
DL-Alpha-Tocopherol	0.10	0.10
Calcium Disodium EDTA	0.07	0.07
Paprika Oleoresin 1000 ASTA	0.02	0.02
Water	396.85	396.85
TOTAL:	1200.00	1200.00

[0028] The ranch dressings were prepared using a POWERGEN homogenizer as the mixing unit. The preservatives, sodium benzoate and potassium sorbate, and EDTA were dissolved in the water. A mixture of the hydrocolloid gums, xanthan gum and propylene glycol alginate, was dispersed in a portion of the vegetable oil (1 part gum in 2 - 5 parts of portion of the oil). The gum dispersion was blended and hydrated in the water at 8,000 rpm for 10 minutes. After blending in the vinegar and lemon juice, the sugar, buttermilk solids, ranch seasoning, salt and titanium dioxide were added and blended at 12,000 rpm for 5 minutes. The polysorbates, 60 and 80 were melted together and blended with the egg yolk in the aqueous phase for about 1 minute at 12,000 rpm.

[0029] Policosanol was added to melted stanol esters and heated in a water-bath to dissolve. The preparation was added to the vegetable oil/CAPROL ET blend at about 140 °F. While mixing the aqueous phase at high speed, 14,000-rpm minimum, the oil preparation was added slowly to the aqueous phase. During the addition of the oil phase, a mixture of the vitamin E acetate, alpha tocopherol and paprika oleoresin was added. After all the oil phase ingredients were added, the dressing emulsion was blended for four more minutes to produce the finished ranch dressing which was filled in bottles and stored. Each dressing contained approximately 10 mg policosanol per 30 gram serving size.

Example 4

[0030] This example discloses cholesterol-lowering Thousand Island dressings containing policosanol with stanol esters and policosanol with sterol esters.

Table 4:

Cholesterol-Lowering Thousand Island Dressings Containing (a) Policosanol with Stanol Esters and (b) Policosanol with Sterol Esters.		
	(a)	(b)
Ingredients	Batch Weight (g)	Batch Weight (g)
Soybean Oil, Salad Oil	17.60	417.60
Relish, Sweet, Ripon	32.00	132.00

Table 4: (continued)

Cholesterol-Lowering Thousand Island Dressings Containing (a) Policosanol with Stanol Esters and (b) Policosanol with Sterol Esters.		
	(a)	(b)
Ingredients	atch Weight (g)	Batch Weight (g)
Sugar, Fine Granulated	26.00	126.00
Vinegar, 120 Grain, White distilled	0.40	80.40
Tomato Paste 31 % Solids	2.00	72.00
Stanol Esters	8.96	
Sterol Esters		68.96
Salt	8.00	18.00
CAPROL ET, A. C. Humko	.36	6.36
Mustard Flour, Baltimore Spice	.00	6.00
Egg Yolk, Liquid, Nepa #2, 10 % Salt	.80	4.80
Polysorbate 60	.60	3.60
KELTROL T Xanthan Gum, Kelco	.12	3.12
Lemon Juice Concentrate, 400 GPL	.40	2.40
Propylene Glycol Alginate, KELCOLOID LVF, Kelco	.56	1.56
Polysorbate 80, TWEEN 80,	.20	1.20
Onion, Minced	.84	0.84
Sodium Benzoate	.60	0.60
Potassium Sorbate	.20	1.20
Garlic Powder	.40	2.40
Black Aquaresin Pepper	.24	0.24
Paprika	.24	0.24
Vitamin E Acetate, Roche	.22	0.22
Policosanol, 95 % Active, Garuda CHOLESTANOL	.42	0.42
Color, Caramel AP 100 Sethness	.12	0.12
DL-Alpha-Tocopherol	.10	0.10
Calcium Disodium EDTA	.07	0.07
Water	49.55	249.55
TOTAL:	200.00	1200.00

[0031] The preservatives (sodium benzoate, potassium sorbate and EDTA) were dissolved in the water. The gums (xanthan and the alginate) were dispersed in a portion of the salad oil (1 part gum in 2 - 5 parts salad oil). Using the POWERGEN homogenizer set at 8,000 rpm, the gum dispersion was blended in the water for 10 minutes. Vinegar and lemon juice were added and blended for 1 minute. Next, the sugar, tomato paste, salt, mustard flour, minced onion and garlic powder were added and blended at 12,000 rpm until the tomato paste was distributed uniformly in the preparation. The polysorbates, 60 and 80 were melted together and blended in the preparation. Liquid egg yolk and caramel color were then added and blended in for about 30 seconds.

[0032] Policosanol was added to the stanol esters. The mixture was heated and melted in a water-bath. The melted policosanol preparation was added to a blend of the salad oil, CAPROL ET, alphas-tocopherol, vitamin E acetate, black aquaresin pepper and paprika oleoresin maintained at about 140 °F. While blending the aqueous phase preparation at about 14,000 rpm, the oil blend was added slowly to form the dressing emulsion. After the oil addition was completed,

the mixing was continued for two more minutes to obtain a tight emulsion. Sweet relish was then mixed gently in the emulsion to complete the process. The dressing was then filled into suitable containers for storage. Each dressing formulation contained approximately 10 mg policosanol per a serving size of 30 grams.

Example 5

[0033] This example describes the application of policosanol dissolved in stanol esters and sterol esters in yogurts. Policosanol was mixed with melted stanol esters or sterol esters. The mixture was heated to approximately 160 - 180 °F in a water bath to melt and dissolve policosanol in the liquid stanol esters or sterol esters. The policosanol preparation was used to produce flavored yogurt products.

Table 5:

Cholesterol-Lowering Yogurt Formulations Containing (a) Policosanol with Stanol Esters and (b) Policosanol with Sterol Esters.		
	(a)	(b)
<u>Yogurt Base Formulation</u>	Batch Weight (g)	Batch Weight (g)
Skim Milk	3560.28	3560.28
Liquid Cane Sugar	451.22	451.212
Condensed Skim Milk	442.50	442.50
Whole Milk (3.4% Milk Fat)	369.00	369.00
Stabilizer Blend (Germantown International)	52.50	52.50
MELOSKIM WP-75 Whey Protein Concentrate (Kerry Foods)	50.00	50.00
Stanol Esters (Raisio)	47.42	-
Sterol Esters (Raisio)	-	47.42
Kosher Edible Gelatin Powder (250) (GMI Products)	16.25	16.25
OPTA-MIST, Modified Food Starch (Opta Foods)	10.00	10.00
Policosanol 95 % Active, Garuda CHOLESTANOL	0.10	0.10
Yogurt Culture YC087 (Chris Hansen)	0.55	0.55
Vitamin E, dl-Alpha-Tocopheryl Acetate (Roche)	0.18	0.18
TOTAL	5000.00	5000.00

Finished Fruit Flavored Yogurt	(a)	(b)
Ingredients	Batch Weight (g)	Batch Weight (g)
Yogurt Base Formulation	4482.00	4482.00
Fruit Preparation	918.00	918.00
TOTAL	5400.00	5400.00
Yogurt serving size, 8 oz (226.8 g) contained ~ 3.6 mg policosanol and ~ 1.0 g stanols or sterols.		

[0034] The laboratory process for preparing the yogurts was as follows. The liquid ingredients including the skim milk, whole milk, condensed milk and liquid sugar were blended together in a stainless steel container in a water-bath on a stove. The starch was dispersed in the liquid blend at medium speed (about 600 rpm). The stove was set at medium heat and the blend was heated slowly. The remaining dry ingredients including gelatin, stabilizers and whey protein concentrate were dispersed into the preparation and the blend was heated to 140 °F. Policosanol was added to stanol ester and heated to a clear liquid, approximately 160 - 180 °F. Vitamin E was added to the policosanol/sterol ester preparation and the mixture was blended with the milk preparation. The blend was heated in the water-bath to 190 °F. After holding the temperature at 190 °F for 5 minutes, the preparation was cooled and homogenized in two stages (1500/500 pounds per square inch) at approximately 140 °F. The homogenized preparation was cooled to about

109 °F and inoculated with the yogurt starter culture. The pH of the yogurt base was monitored hourly until pH 4.5 was attained. The preparation was agitated and cooled to 65 °F and stored overnight in a refrigerator. The finished yogurt was packed in suitable containers after mixing 83 parts of the yogurt base with 17 parts fruit flavor preparations.

5 Example 6

[0035] This example discloses the application of policosanol dissolved in stanol esters or sterol esters in cholesterol-lowering pasteurized cheese spread formulation.

10 Table 6:

Cholesterol-Lowering Pasteurized Cheese Spread Formulations Containing (a) Policosanol and Stanol Esters and (b) Policosanol and Sterol Esters.		
	(a)	(b)
Ingredients	Batch Size (g)	Batch Size (g)
Light Cheddar Cheese (Cabot)	1275.00	1275.00
Stanol Esters (Raisio)	287.50	-
Sterol Esters (Raisio)	-	287.50
Policosanol, 95 % Active, Garuda CHOLESTANOL	0.62	0.62
JOHA S-4, Sodium Phosphates (B K Ladenburg)	50.00	50.00
JOHA S-9 (Sodium Phosphates (B K Ladenburg)	12.50	12.50
Sodium Citrate	6.25	6.25
Nonfat Dry Milk Solids	62.50	62.50
Lactic Acid	10.00	10.00
Salt	20.82	20.82
Annatto Color (Chris Hansen)	0.42	0.42
Water	774.39	774.39
Total	2500.00	2500.00

[0036] The laboratory process for preparing the cheese spread was as follows. The cheddar cheese was weighed in a stainless steel container. The container was heated in a water-bath on a stove to melt the cheese. Policosanol was added to stanol esters and heated in a water-bath to a clear liquid. The liquid policosanol preparation was transferred and blended with the melted cheddar cheese. About half the volume of the water was added and blended with the cheese preparation to a uniform consistency. The salt, nonfat dry milk solids, phosphates, citrate, lactic acid, annatto color and remaining water were added and blended to uniform consistency. The temperature of the cheese preparation was raised and held at 176 °F for 5 minutes. The finished product was packed hot into suitable containers and stored refrigerated. Each formulation contained approximately 3.5 mg of policosanol per 15 g of the processed cheese spread.

45 Example 7

[0037] This example concerns the application of policosanol dissolved in stanol esters or sterol esters in preparing a cholesterol-lowering processed cheese sauce.

50 Table 7:

Cholesterol-Lowering Processed Cheese Sauce Containing (a) Policosanol and Stanol Esters and (b) Policosanol and Sterol Esters.		
	(a)	(b)
Ingredients	Batch Weight (g)	Batch Weight (g)
Light Cheddar Cheese (Cabot)	1075.00	1075.00

Table 7: (continued)

Cholesterol-Lowering Processed Cheese Sauce Containing (a) Policosanol and Stanol Esters and (b) Policosanol and Sterol Esters.		
	(a)	(b)
Ingredients	Batch Weight (g)	Batch Weight (g)
Stanol Esters (Raisio)	287.50	-
Sterol Esters (Raisio)	-	287.50
Policosanols, 95 % Active, Garuda CHOLESTANOL	0.62	0.62
Sodium Citrate	62.50	62.50
Nonfat Dry Milk Solids	83.25	83.25
Lactic Acid	10.00	10.00
Salt	25.00	25.00
Annatto Color (Chris Hansen)	0.42	0.42
Water	955.71	955.71
Total	2500.00	2500.00

[0038] The cheddar cheese sauce was prepared as follows. The cheddar cheese was melted in a suitable container placed in a water-bath heated on a stove. Policosanol was added to the stanol or sterol esters. The container was heated to about 160 - 180 °F in a water-bath to melt and dissolve policosanol in the stanol or sterol esters. The liquid policosanol preparation was blended with the melted cheese. About half the volume of the water was added to the melted cheese and blended to uniform consistency. After adding and mixing in the salt, nonfat milk solids, citrate, lactic acid and annatto color, the remaining water was added and blended to uniform consistency. The cheese sauce was then heated and the temperature held for five minutes at 176 °F. The finished cheese sauce was packed hot in suitable containers and stored refrigerated. Each formulation contained approximately 3.5 mg of policosanol per 15 g of the processed cheese sauce.

Example 8

[0039] This example discloses the application of policosanol dissolved in stanol or stanol esters in typical cholesterol-lowering nutritional bars.

Table 8:

Cholesterol-Lowering Chocolate Coated Nutritional Bar Formulations Containing (a) Policosanol and Stanol Esters and (b) Policosanol and Sterol Esters.		
	(a)	(b)
Ingredients	Batch Weight (g)	Batch Weight (g)
Corn Syrup, 63 Dextrose Equivalents (Cargill)	390.36	390.36
KRYSTAR Crystalline Sugar (Far Best Foods)	70.20	70.20
CENTROPHASE C Lecithin (Central Soya)	4.80	4.80
Stanol Esters	61.44	-
Sterol Esters	-	61.44
Policosanols, 95 % Active, Garuda CHOLESTANOL	0.13	0.13
Crisp Rice	99.18	99.18
Cocoa Powder (Paragon)	76.00	76.00
Oats, Baby	73.40	73.40

Table 8: (continued)

Cholesterol-Lowering Chocolate Coated Nutritional Bar Formulations Containing (a) Policosanol and Stanol Esters and (b) Policosanol and Sterol Esters.		
	(a)	(b)
Ingredients	Batch Weight (g)	Batch Weight (g)
Soy Flour, Nutrisoy	73.40	73.40
Centex Textured Soy Flour, Unfortified (Central Soya)	64.00	64.00
Rice Flour, Pregelatinized	26.46	26.46
Vitamin E, 25 % RDA Blend	12.14	12.14
Ricex Rice Bran	9.60	9.60
Chocolate Fudge Flavor	9.12	9.12
Chocolate Flavor	4.74	4.74
Caramel Flavor	1.09	1.09
Target Coating	223.94	223.94
Total	1200.00	1200.00

[0040] The corn syrup, sugar and lecithin were blended together in a stainless steel container and heated gently in a water-bath on a stove. The stanol esters and policosanol were mixed together and heated to melt at about 160 - 180 °F. The clear liquid policosanol preparation was blended gently into the hot syrup at about 160 °F. Vitamin E, the chocolate fudge, chocolate and caramel flavors were blended in the syrup preparation. The syrup preparation was transferred into a bench top Hobart bowl with a dough spindle mixer. The remaining dry ingredients were mixed together in a plastic container. The mixture was mixed gently at speed 1 in the syrup preparation to a uniform consistency. The dough was transferred into a pan and rolled with a wooden roller to desired thickness. The dough was covered with a wax paper and cooled in a refrigerator to firm up before cutting it into pieces. The pieces were then enrobed with the target coating and chilled through a cooling tunnel. The finished nutritional bars were then wrapped in foil and heat-sealed. Each formulation contained approximately, 3.5 mg of policosanol per 34 g serving size of the nutritional bar.

Claims

1. A comestible comprising:
an effective amount of a long chain alcohol or mixture of long chain alcohols sufficient to lower cholesterol in a vertebrate, said long chain alcohol or alcohols being admixed with an effective amount of a sterol compound sufficient to lower cholesterol in a vertebrate, said long chain alcohol/sterol composition admixture being provided in a foodstuff.
2. The comestible of claim 1 wherein the amount of long chain alcohol is from 0.1 to 40 milligrams per serving, preferably from 1.0 to 20 milligrams per serving.
3. The comestible of claim 1 or claim 2 wherein the long chain alcohol is predominately octacosanol providing about 10 milligrams of octacosanol per serving.
4. The comestible of any one of claims 1 to 3 wherein the sterol compound is provided in an amount from 0.4 to 4.0 grams per serving.
5. The comestible of any one of claims 1 to 4 wherein the sterol compound is a sterol, stanol, sterol ester, stanol ester or oryzanol.
6. The comestible of claim 1 which is an oil, such as a cooking oil, a margarine or a salad dressing.

7. A method of preparing a comestible comprising:

providing a long chain alcohol in an amount sufficient to lower cholesterol in human;
providing a sterol compound in an amount sufficient to lower cholesterol in a human;
5 admixing said long chain alcohol and said sterol compound; and adding said admixture to a comestible.

8. The method of claim 7 wherein the long chain alcohol and sterol are heated.

9. The method of claim 7 or claim 8 wherein the admixture additionally contains a surfactant and/or an emulsifying
10 agent.

10. The method of any one of claims 7 to 9 wherein the long chain alcohol and sterol admixture forms an emulsion.

11. The comestible of any one of claims 1 to 6 which contains less than about 40 mg of a long chain alcohol and 0.4
15 to 4 grams of a sterol for use in treating hypercholesterolemia or for reducing cholesterol in a vertebrate.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 108 364 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
29.05.2002 Bulletin 2002/22

(51) Int Cl.7: **A23D 9/007**, **A23L 1/24**,
A23L 1/30

(43) Date of publication A2:
20.06.2001 Bulletin 2001/25

(21) Application number: **00311172.1**

(22) Date of filing: **14.12.2000**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• **Dartey, Clemence K.**
North Wales, PA 19545 (US)
• **Sox, Thomas E.**
Ambler, PA 19002 (US)

(30) Priority: **15.12.1999 US 461607**

(71) Applicant: **McNEIL-PPC, INC.**
Skillman, NJ 08858 (US)

(74) Representative: **Mercer, Christopher Paul**
Carpmaels & Ransford
43, Bloomsbury Square
London WC1A 2RA (GB)

(54) **Long chain alcohols admixed in sterol compounds**

(57) The present invention provides a method of providing a desirable admixture of long chain alcohols and sterols in a food or beverage product. More specifically, this process comprises dissolving or suspending

the long chain alcohol in the sterol, and then incorporating the admixture into food or beverage product.

EP 1 108 364 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 31 1172

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X,D	US 5 952 393 A (SORKIN JR HARLAN LEE) 14 September 1999 (1999-09-14) * column 1, line 5 - line 8 * * column 2, line 26 - line 33 * * column 3, line 13 - line 26; claims 1-10; example 2 *	1-5,11	A23D9/007 A23L1/24 A23L1/30
X	US 3 865 939 A (JANDACEK RONALD JAMES) 11 February 1975 (1975-02-11) * column 1, line 42 - line 64 * * column 3, line 12 - line 28 * * column 6, line 13 - line 20; claims 1-4,7; example 1 *	1,2,4-7	
X	US 5 244 887 A (STRAUB CARL D) 14 September 1993 (1993-09-14) * column 5, line 56 - column 6, line 22 * * column 6, line 45 - column 7, line 17; claims 1,2 *	1,2,4-7	
X	DATABASE WPI Section Ch, Week 198724 Derwent Publications Ltd., London, GB; Class A96, AN 1987-167050 XP002190062 & JP 62 099323 A (HAGIWARA Y), 8 May 1987 (1987-05-08) * abstract *	1,2,5	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A23D A23C A23L A23P A61K
X	WO 96 10033 A (UNIV BRITISH COLUMBIA) 4 April 1996 (1996-04-04) * page 1, line 9 - page 5, line 9 * * page 10, line 11 - line 12 * * page 11, line 20 - line 27; claims 8-18; examples 2-7 *	1-5,11	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 27 March 2002	Examiner Muller, I
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03 82 (Pst/C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 31 1172

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	WO 98 47385 A (HAILES ANNE ; UNILEVER PLC (GB); VERMAAS LEO FRANS (NL); ZWEMMER OL) 29 October 1998 (1998-10-29) * page 1 - page 2; claims 1-11; example I; table I *	1-11	
Y	WO 96 38047 A (UNILEVER PLC ; UNILEVER NV (NL); LIEVENSE LOURUS CORNELIS (NL)) 5 December 1996 (1996-12-05) * page 1 - page 3; claims 1-7; examples I-VII *	1-11	
E	WO 01 32031 A (PIRAKITIKULR PICHAI ; MENON VINOD P (US); MONSANTO CO (US); KINLEN) 10 May 2001 (2001-05-10) * page 6, line 8 - page 9, line 4 * * page 10, line 29 - line 31 * * page 21, line 15 - page 22, line 11; claims 5,16,29,44-49; example 1 *	1-5,7,9	
E	WO 01 32036 A (MENON VINOD P ; MONSANTO CO (US); KINLEN PATRICK J (US); PIRAKITIKU) 10 May 2001 (2001-05-10) * claims 44-49; example 11 *	1-5	
E	EP 1 177 729 A (HAERTING THOMAS FRANCIS) 6 February 2002 (2002-02-06) * page 2, line 1 - line 21 * * page 3, line 16 - line 20 * * page 4, line 5 - line 12; claims 1-9; examples 4,6,7,9,11 *	1,2,4-11	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 27 March 2002	Examiner Muller, I
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 82 (P4/C21)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 31 1172

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

27-03-2002

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
US 5952393	A	14-09-1999	AU	3290299 A	30-08-1999
			CA	2321750 A1	19-08-1999
			CN	1294515 T	09-05-2001
			EP	1061927 A1	27-12-2000
			JP	2002502879 T	29-01-2002
			WO	9940922 A1	19-08-1999
US 3865939	A	11-02-1975	BE	811452 A1	22-08-1974
			CA	1024814 A1	24-01-1978
			DE	2408067 A1	05-09-1974
			FR	2218838 A1	20-09-1974
			GB	1413102 A	05-11-1975
			IT	1047132 B	10-09-1980
			JP	1135221 C	14-02-1983
			JP	50040605 A	14-04-1975
			JP	57026732 B	07-06-1982
			NL	7402426 A ,C	27-08-1974
US 5244887	A	14-09-1993	NONE		
JP 62099323	A	08-05-1987	JP	1818104 C	27-01-1994
			JP	5029206 B	28-04-1993
WO 9610033	A	04-04-1996	AU	3559795 A	19-04-1996
			BG	101354 A	31-03-1998
			BR	9509119 A	28-10-1997
			CA	2201328 A1	04-04-1996
			WO	9610033 A1	04-04-1996
			CN	1166174 A ,B	26-11-1997
			CZ	9700927 A3	18-02-1998
			EE	9700073 A	15-10-1997
			EP	1148062 A2	24-10-2001
			EP	0783514 A1	16-07-1997
			HU	77522 A2	28-05-1998
			JP	10506394 T	23-06-1998
			LT	97074 A ,B	27-10-1997
			LV	11900 A	20-12-1997
			LV	11900 B	20-05-1998
			MD	970178 A	28-02-1999
			NO	971304 A	14-05-1997
			NO	11062 A	14-05-1997
			PL	319384 A1	04-08-1997
			RU	2165431 C2	20-04-2001
			SK	41597 A3	05-11-1997
			US	5770749 A	23-06-1998

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 31 1172

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

27-03-2002

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9847385	A	29-10-1998	AU	7430698 A	13-11-1998
			WO	9847385 A1	29-10-1998
			EP	0977489 A1	09-02-2000
			US	6277430 B1	21-08-2001
WO 9638047	A	05-12-1996	AU	1347500 A	20-04-2000
			AU	1347600 A	20-04-2000
			AU	6003796 A	18-12-1996
			BR	9608914 A	02-03-1999
			CA	2222771 A1	05-12-1996
			DE	69606638 D1	16-03-2000
			DE	69606638 T2	20-07-2000
			DK	828434 T3	13-06-2000
			WO	9638047 A1	05-12-1996
			EP	0828434 A1	18-03-1998
			EP	0960567 A2	01-12-1999
			EP	0962150 A2	08-12-1999
			ES	2142589 T3	16-04-2000
			JP	11506324 T	08-06-1999
			ZA	9604554 A	03-12-1997
			AU	1347200 A	06-04-2000
			AU	1347300 A	06-04-2000
			AU	1347400 A	06-04-2000
WO 0132031	A	10-05-2001	AU	1460801 A	14-05-2001
			WO	0132031 A2	10-05-2001
WO 0132036	A	10-05-2001	AU	1463601 A	14-05-2001
			WO	0132036 A1	10-05-2001
EP 1177729	A	06-02-2002	EP	1177729 A2	06-02-2002